

Baryon asymmetry from hypermagnetic helicity in dilaton hypercharge electromagnetism

Kazuharu Bamba^{* †}

*Yukawa Institute for Theoretical Physics,
Kyoto University, Kyoto 606-8502, Japan*

Abstract

The generation of the baryon asymmetry of the Universe (BAU) from the hypermagnetic helicity, the physical interpretation of which is given in terms of hypermagnetic knots, is studied in inflationary cosmology, taking into account the breaking of the conformal invariance of hypercharge electromagnetic fields through both a coupling with the dilaton and that with a pseudoscalar field. It is shown that if the electroweak phase transition (EWPT) is strongly first order and the present amplitude of the generated magnetic fields on the horizon scale is sufficiently large, a baryon asymmetry with a sufficient magnitude to account for the observed baryon to entropy ratio can be generated.

PACS numbers: 98.80.Cq, 11.30.Fs, 98.62.En

^{*} Research Fellow of the Japan Society for the Promotion of Science

[†] E-mail: bamba@yukawa.kyoto-u.ac.jp

I. INTRODUCTION

It is observationally known that there exists a net excess of baryons over antibaryons in the Universe. In order to generate the baryon asymmetry of the Universe (BAU) from a state in which baryons and antibaryons have same abundances, the Sakharov's three conditions [1] must be satisfied: (1) baryon number nonconservation, (2) C and CP violation, and (3) a departure from thermal equilibrium. Many cosmological scenarios in which the above three conditions could be satisfied have been proposed (for reviews of the BAU, see Refs. [2, 3]). The origin of the BAU, however, is not well established yet.

It has been pointed out [4, 5, 6, 7, 8] that hypercharge electromagnetic fields could play a significant role in the electroweak (EW) scenario [9, 10, 11, 12] for baryogenesis. Giovannini and Shaposhnikov [5, 6] have shown that the Chern-Simons number stored in the hypercharge electromagnetic fields, i.e., the hypermagnetic helicity, is converted into fermions at the electroweak phase transition (EWPT) owing to the Abelian anomaly [13], and at the same time the hypermagnetic fields are replaced by the ordinary magnetic fields, which survive after the EWPT. The hypermagnetic helicity corresponds to a topologically non-trivial hypercharge configurations. Topologically non-trivial refers here to the topology of the magnetic flux lines. The physical interpretation of these hypercharge configurations is given in terms of hypermagnetic knots. The generated fermions will not be destroyed by sphaleron processes [14] if the EWPT is strongly first order. This condition could be met in the extensions of the minimal standard model (MSM) [15].

The most natural origin of large-scale hypermagnetic fields before the EWPT is hypercharge electromagnetic quantum fluctuations generated in the inflationary stage [16] (for reviews of inflation, see Refs. [17, 18]). This is because inflation naturally produces effects on very large scales, larger than Hubble horizon, starting from microphysical processes operating on a causally connected volume. If the conformal invariance of the Maxwell theory is broken by some mechanism in the inflationary stage, hypercharge electromagnetic quantum fluctuations could be generated even in the Friedmann-Robertson-Walker (FRW) spacetime, which is conformally flat. Hence several breaking mechanisms have been proposed in Refs. [16, 19, 20, 21, 22, 23, 24, 25] (for other mechanisms, see reviews of the origin of cosmic magnetic fields [26, 27, 28, 29, 30] and references therein). In particular, it follows from indications in higher-dimensional theories including string theory that there can exist the dilaton field coupled to the hypercharge electromagnetic fields. This coupling also breaks the conformal invariance of the Maxwell theory [19, 20, 24, 25] and hence the large-scale hypercharge magnetic fields could be generated.

Furthermore, it has been noticed [31, 32] that when a pseudoscalar field ϕ with an axion-like coupling to the $U(1)_Y$ hypercharge field strength $Y_{\mu\nu}$ in the form $\phi Y_{\mu\nu} \tilde{Y}^{\mu\nu}$, where $\tilde{Y}^{\mu\nu}$ is the dual of $Y_{\mu\nu}$ and $Y_{\mu\nu} \tilde{Y}^{\mu\nu}$ corresponds to the hypercharge topological number density, coherently rolls or oscillates before the EWPT, the Sakharov's three conditions can be satisfied. In this case, the motion of the pseudoscalar field generates a time-dependent hypercharge topological number condensate which violates fermion number conservation through the Abelian anomalous coupling [13], and realizes a departure from equilibrium. Moreover, C symmetry is violated by the chiral coupling of the hypercharge gauge boson to fermions. Furthermore, since the hypercharge topological number is odd under CP , CP symmetry is spontaneously broken if the hypercharge topological number is coupled to a field with a time-dependent expectation value. This mechanism is able to generate a net Chern-Simons number which can survive until the EWPT and then be converted into a

baryon asymmetry. Pseudoscalar fields with the above axion-like coupling appear in several possible extensions of the standard model [33]. Incidentally, the generation of large-scale magnetic fields owing to the breaking of the conformal invariance of the Maxwell theory through such an axion-like coupling between a pseudoscalar field and electromagnetic fields have been considered in Refs. [16, 21, 22]. Moreover, baryogenesis due to the above coupling has been discussed in Ref. [34]. Furthermore, the generation of the magnetic helicity owing to the above coupling has been considered in Ref. [35].

In the present paper, in addition to the existence of the dilaton coupled to hypercharge electromagnetic fields, we assume the existence of a pseudoscalar field with an axion-like coupling to hypercharge electromagnetic fields, and consider the generation of the BAU in slow-roll exponential inflation models. In particular, we consider the generation of the Chern-Simons number, i.e., the hypermagnetic helicity, through the coupling between a pseudoscalar field and hypercharge electromagnetic fields in the inflationary stage. The generated Chern-Simons number is converted into fermions at the EWPT owing to the Abelian anomaly, and the generated fermions can survive after the EWPT if the EWPT is strongly first order. The reason why we consider the generation of the Chern-Simons number during inflation is as follows: In this scenario, the BAU is induced by the helicity of the hypermagnetic fields. Hence the large-scale hypermagnetic fields whose present scale is larger than the present horizon scale have to be generated in order that the resultant BAU could be homogeneous over the present horizon.

This paper is organized as follows. In Sec. II we describe our model and derive equations of motion from its action. In Sec. III we consider the evolution of the $U(1)_Y$ gauge field and investigate the generated Chern-Simons number density, and then estimate the resultant baryon asymmetry. Finally, Sec. IV is devoted to a conclusion.

We use units in which $k_B = c = \hbar = 1$ and denote the gravitational constant $8\pi G$ by κ^2 so that $\kappa^2 \equiv 8\pi/M_{\text{Pl}}^2$ where $M_{\text{Pl}} = G^{-1/2} = 1.2 \times 10^{19} \text{GeV}$ is the Planck mass. Moreover, in terms of electromagnetism we adopt Heaviside-Lorentz units. Throughout the present paper, the subscripts ‘1’, ‘R’, and ‘0’ represent the quantities at the time t_1 when a given mode of the $U(1)_Y$ gauge field first crosses outside the horizon during inflation, the end of inflation (namely, the instantaneous reheating stage) t_R , and the present time t_0 , respectively.

II. MODEL

A. Action

We introduce the dilaton field Φ and a pseudoscalar field ϕ . Moreover, we introduce the coupling of the dilaton to hypercharge electromagnetic fields and that of the pseudoscalar to those fields. Our model action is the following:

$$S = \int d^4x \sqrt{-g} [\mathcal{L}_{\text{inflaton}} + \mathcal{L}_{\text{dilaton}} + \mathcal{L}_{\text{ps}} + \mathcal{L}_{\text{HEM}}], \quad (2.1)$$

$$\mathcal{L}_{\text{inflaton}} = -\frac{1}{2} g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi - U[\varphi], \quad (2.2)$$

$$\mathcal{L}_{\text{dilaton}} = -\frac{1}{2} g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi - V[\Phi], \quad (2.3)$$

$$\mathcal{L}_{\text{ps}} = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - W[\phi], \quad (2.4)$$

$$\mathcal{L}_{\text{HEM}} = -\frac{1}{4}f(\Phi)\left(Y_{\mu\nu}Y^{\mu\nu} + g_{\text{ps}}\frac{\phi}{M}Y_{\mu\nu}\tilde{Y}^{\mu\nu}\right), \quad (2.5)$$

$$f(\Phi) = \exp(-\lambda\kappa\Phi), \quad (2.6)$$

$$V[\Phi] = \bar{V}\exp(-\tilde{\lambda}\kappa\Phi), \quad (2.7)$$

$$W[\phi] = \frac{1}{2}m^2\phi^2, \quad (2.8)$$

where g is the determinant of the metric tensor $g_{\mu\nu}$, $U[\varphi]$, $V[\Phi]$, and $W[\phi]$ are the inflaton, the dilaton, and the pseudoscalar potentials, respectively, $\bar{V}$ is a constant, m is the mass of the pseudoscalar field, and f is the coupling between the dilaton and hypercharge electromagnetic fields with λ^1 and $\tilde{\lambda}$ (> 0) being dimensionless constants. Moreover, $g_{\text{ps}} = \bar{g}_{\text{ps}}\alpha'/(2\pi)$, where $\bar{g}_{\text{ps}}$ is a numerical factor and $\alpha' = g'^2/(4\pi)$. Here, g' is the $U(1)_Y$ gauge coupling constant. M denotes a mass scale. Furthermore, the $U(1)_Y$ hypercharge field strength is given by $Y_{\mu\nu} = \nabla_\mu Y_\nu - \nabla_\nu Y_\mu$, where $\tilde{Y}^{\mu\nu} = (1/2)\epsilon^{\mu\nu\rho\sigma}Y_{\rho\sigma}$. Here, Y_μ is the $U(1)_Y$ gauge field, ∇_μ denotes the covariant derivative, and $\epsilon^{\mu\nu\rho\sigma}$ is the Levi-Civita tensor. Moreover, we note that covariant derivatives for the antisymmetric tensor $Y_{\mu\nu}$ in the metric (2.13) as is shown in the next subsection are simple derivatives, $\nabla_\mu = \partial_\mu$, as it appears later in the equations of motion (2.12) and (2.22) based on the hypercharge electromagnetic part of our model Lagrangian in Eq. (2.5).

B. Equations of motion

From the above action in Eq. (2.1), the equations of motion for the inflaton, the dilaton, the pseudoscalar field, and hypercharge electromagnetic fields can be derived as follows:

$$-\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\varphi) + \frac{dU[\varphi]}{d\varphi} = 0, \quad (2.9)$$

$$-\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\Phi) + \frac{dV[\Phi]}{d\Phi} = -\frac{1}{4}\frac{df(\Phi)}{d\Phi}\left(Y_{\mu\nu}Y^{\mu\nu} + g_{\text{ps}}\frac{\phi}{M}Y_{\mu\nu}\tilde{Y}^{\mu\nu}\right), \quad (2.10)$$

$$-\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\phi) + \frac{dW[\phi]}{d\phi} = -\frac{1}{4}\frac{g_{\text{ps}}}{M}f(\Phi)Y_{\mu\nu}\tilde{Y}^{\mu\nu}, \quad (2.11)$$

$$\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}f(\Phi)Y^{\mu\nu}) = -\frac{g_{\text{ps}}}{M}\partial_\mu[f(\Phi)\phi]\tilde{Y}^{\mu\nu}. \quad (2.12)$$

We now assume the spatially flat Friedmann-Robertson-Walker (FRW) space-time with

¹ The sign of λ in the present paper is opposite to that in Refs. [24, 25]

the metric

$$\begin{aligned} ds^2 &= g_{\mu\nu} dx^\mu dx^\nu = -dt^2 + a^2(t) d\mathbf{x}^2 \\ &= a^2(\eta)(-d\eta^2 + d\mathbf{x}^2), \end{aligned} \quad (2.13)$$

where a is the scale factor, and η is the conformal time.

Since we are interested in the specific case in which the background space-time is inflating, we assume that the spatial derivatives of φ , ϕ , and Φ are negligible compared to the other terms (if this is not the case at the beginning of inflation, any spatial inhomogeneities will quickly be inflated away and this assumption will quickly become very accurate). Hence the equations of motion for the background homogeneous scalar fields read

$$\ddot{\varphi} + 3H\dot{\varphi} + \frac{dU[\varphi]}{d\varphi} = 0, \quad (2.14)$$

$$\ddot{\Phi} + 3H\dot{\Phi} + \frac{dV[\Phi]}{d\Phi} = 0, \quad (2.15)$$

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dW[\phi]}{d\phi} = 0, \quad (2.16)$$

together with the background Friedmann equation

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa^2}{3}(\rho_\varphi + \rho_\Phi + \rho_\phi), \quad (2.17)$$

$$\rho_\varphi = \frac{1}{2}\dot{\varphi}^2 + U[\varphi], \quad (2.18)$$

$$\rho_\Phi = \frac{1}{2}\dot{\Phi}^2 + V[\Phi], \quad (2.19)$$

$$\rho_\phi = \frac{1}{2}\dot{\phi}^2 + W[\phi], \quad (2.20)$$

where a dot denotes a time derivative. Here, ρ_φ , ρ_Φ , and ρ_ϕ are the energy densities of the inflaton, the dilaton, and the pseudoscalar field, respectively. We here consider the case in which slow-roll exponential inflation is driven by the potential energy of the inflaton and during inflation the energy densities of the dilaton and the pseudoscalar field are much smaller than that of the inflaton, $\rho_\varphi \gg \rho_\Phi$, $\rho_\varphi \gg \rho_\phi$. Hence, during inflation H reads

$$H^2 \approx \frac{\kappa^2}{3}\rho_\varphi \equiv H_{\text{inf}}^2, \quad (2.21)$$

where H_{inf} is the Hubble constant in the inflationary stage.

We consider the evolution of the $U(1)_Y$ gauge field in this background. Its equation of motion in the Coulomb gauge, $Y_0(t, \mathbf{x}) = 0$ and $\partial_j Y^j(t, \mathbf{x}) = 0$, becomes

$$\ddot{Y}_i(t, \mathbf{x}) + \left(H + \frac{\dot{f}}{f}\right) \dot{Y}_i(t, \mathbf{x}) - \frac{1}{a^2} \partial_j \partial_j Y_i(t, \mathbf{x}) - \frac{g_{\text{ps}}}{M} \frac{1}{af} \frac{d(f\phi)}{dt} \epsilon^{ijk} \partial_j Y_k(t, \mathbf{x}) = 0. \quad (2.22)$$

III. GENERATION OF THE BAU

In this section, we consider the generation of the BAU. First, we consider the evolution of the dilaton, that of the pseudoscalar field, and that of the $U(1)_Y$ gauge field. Next, we investigate the generation of the Chern-Simons number density in the inflationary stage, and then estimate the ratio of the baryonic number density to the entropy density.

A. Evolution of the dilaton and that of the pseudoscalar field

In this subsection, we consider the evolution of the dilaton and the pseudoscalar field. Here we consider the case in which slow-roll exponential inflation is realized and the scale factor $a(t)$ is given by

$$a(t) = a_1 \exp [H_{\text{inf}}(t - t_1)], \quad (3.1)$$

where a_1 is the scale factor at the time t_1 when a given comoving wavelength $2\pi/k$ of the $U(1)_Y$ gauge field first crosses outside the horizon during inflation, $k/(a_1 H_{\text{inf}}) = 1$.

First, we investigate the evolution of the dilaton. We consider the case in which we can apply slow-roll approximation to the dilaton, that is,

$$\left| \frac{\ddot{\Phi}}{H_{\text{inf}} \dot{\Phi}} \right| \ll 1, \quad (3.2)$$

and then Eq. (2.15) is reduced to

$$3H_{\text{inf}} \dot{\Phi} + \frac{dV[\Phi]}{d\Phi} = 0. \quad (3.3)$$

The solution of this equation is given by

$$\Phi = \frac{1}{\tilde{\lambda}\kappa} \ln \left[\tilde{\lambda}^2 w H_{\text{inf}} (t - t_R) + \exp \left(\tilde{\lambda} \kappa \Phi_R \right) \right], \quad (3.4)$$

$$w \equiv \frac{\bar{V}}{3H_{\text{inf}}^2/\kappa^2} \approx \frac{\bar{V}}{\rho_\varphi}, \quad (3.5)$$

where Φ_R is the dilaton field amplitude at the end of inflation. Here we consider the case in which after inflation, the dilaton is finally stabilized when it feels other contributions to its potential, e.g., from gaugino condensation [36] that generates a potential minimum [37, 38]. As it reaches there, the dilaton starts oscillation and finally decays into radiation at $t = t_R$. Hence we assume that the potential minimum is generated at $\Phi = \Phi_R = 0$, so that the coupling f between the dilaton and hypercharge electromagnetic fields is set to unity and thus the standard Maxwell theory is recovered. Moreover, in deriving the second approximate equality in Eq. (3.5), we have used Eq. (2.21). Since we have $\rho_\varphi \gg \rho_\Phi$ by assumption, $w \ll 1$.

It follows from Eq. (3.4) that the slow-roll condition to the dilaton, Eq. (3.2), is equivalent to the following relation:

$$\left| \frac{\ddot{\Phi}}{H_{\text{inf}} \dot{\Phi}} \right| \ll 1 \iff \tilde{\lambda}^2 \frac{V[\Phi]}{\rho_\varphi} \ll 1. \quad (3.6)$$

In deriving this relation, we have used Eq. (2.21). If we assume that $\tilde{\lambda} \sim \mathcal{O}(1)$, the second relation in (3.6), $\tilde{\lambda}^2 V[\Phi]/\rho_\varphi \ll 1$, is satisfied during inflation because $\rho_\varphi \gg \rho_\Phi$.

Next, we consider the evolution of the pseudoscalar field. The solution of Eq. (2.16) with Eq. (2.8) is given by [21]

$$\phi = \phi_1 \exp \left\{ \frac{3}{2} \left[-1 \pm \sqrt{1 - \left(\frac{2m}{3H_{\text{inf}}} \right)^2} \right] H_{\text{inf}} (t - t_1) \right\}. \quad (3.7)$$

For $m \gg H_{\text{inf}}$, we find that the approximate solution of Eq. (2.16) with Eq. (2.8) is given by [21]

$$\phi \approx \phi_1 \exp \left[-\frac{3}{2} H_{\text{inf}} (t - t_1) \right] \sin \left[m (t - t_1) + \frac{\pi}{2} \right]. \quad (3.8)$$

In deriving Eq. (3.8), we have used Eq. (3.1). Moreover, we consider the case in which after inflation, the pseudoscalar field ϕ decays in the radiation-dominated stage and hence the entropy per comoving volume remains practically constant.

B. Evolution of the $U(1)_Y$ gauge field

Next, we consider the evolution of the $U(1)_Y$ gauge field. To begin with, we shall quantize the $U(1)_Y$ gauge field $Y_\mu(t, \mathbf{x})$. It follows from the hypercharge electromagnetic part of our model Lagrangian in Eq. (2.5) that the canonical momenta conjugate to $Y_\mu(t, \mathbf{x})$ are given by

$$\pi_0 = 0, \quad \pi_i = f(\Phi) a(t) \dot{Y}_i(t, \mathbf{x}). \quad (3.9)$$

We impose the canonical commutation relation between $Y_i(t, \mathbf{x})$ and $\pi_j(t, \mathbf{x})$,

$$[Y_i(t, \mathbf{x}), \pi_j(t, \mathbf{y})] = i \int \frac{d^3 k}{(2\pi)^3} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right), \quad (3.10)$$

where $\mathbf{k}$ is comoving wave number, and k denotes its amplitude $|\mathbf{k}|$. From this relation, we obtain the expression for $Y_i(t, \mathbf{x})$ as

$$Y_i(t, \mathbf{x}) = \int \frac{d^3 k}{(2\pi)^{3/2}} \left[\hat{b}(\mathbf{k}) Y_i(t, \mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} + \hat{b}^\dagger(\mathbf{k}) Y_i^*(t, \mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right], \quad (3.11)$$

where $\hat{b}(\mathbf{k})$ and $\hat{b}^\dagger(\mathbf{k})$ are the annihilation and creation operators which satisfy

$$[\hat{b}(\mathbf{k}), \hat{b}^\dagger(\mathbf{k}')] = \delta^3(\mathbf{k} - \mathbf{k}'), \quad [\hat{b}(\mathbf{k}), \hat{b}(\mathbf{k}')] = [\hat{b}^\dagger(\mathbf{k}), \hat{b}^\dagger(\mathbf{k}')] = 0. \quad (3.12)$$

It follows from Eqs. (3.10) and (3.11) that the normalization condition for $Y_i(k, t)$ reads

$$Y_i(k, t) \dot{Y}_j^*(k, t) - \dot{Y}_j(k, t) Y_i^*(k, t) = \frac{i}{fa} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right). \quad (3.13)$$

From now on we choose the x^3 axis to lie along the spatial momentum direction $\mathbf{k}$ and denote the transverse directions x^I with $I = 1, 2$. From Eq. (2.22), we find that the Fourier modes $Y_I(k, t)$ of the $U(1)_Y$ gauge field satisfy the following equations:

$$\ddot{Y}_1(k, t) + \left(H_{\text{inf}} + \frac{\dot{f}}{f}\right) \dot{Y}_1(k, t) + \frac{k^2}{a^2} Y_1(k, t) + ik \frac{g_{\text{ps}}}{M} \frac{1}{af} \frac{d(f\phi)}{dt} Y_2(k, t) = 0, \quad (3.14)$$

$$\ddot{Y}_2(k, t) + \left(H_{\text{inf}} + \frac{\dot{f}}{f}\right) \dot{Y}_2(k, t) + \frac{k^2}{a^2} Y_2(k, t) - ik \frac{g_{\text{ps}}}{M} \frac{1}{af} \frac{d(f\phi)}{dt} Y_1(k, t) = 0. \quad (3.15)$$

In order to decouple the system of Eqs. (3.14) and (3.15), we consider circular polarizations expressed by the combination of linear polarizations as $Y_{\pm}(k, t) \equiv Y_1(k, t) \pm iY_2(k, t)$. From Eqs. (3.14) and (3.15), we find that $Y_{\pm}(k, t)$ satisfies the following equation:

$$\ddot{Y}_{\pm}(k, t) + \left(H_{\text{inf}} + \frac{\dot{f}}{f}\right) \dot{Y}_{\pm}(k, t) + \left[\left(\frac{k}{a}\right)^2 \pm \frac{g_{\text{ps}}}{M} \left(\frac{\dot{f}}{f}\phi + \dot{\phi}\right) \left(\frac{k}{a}\right)\right] Y_{\pm}(k, t) = 0. \quad (3.16)$$

Since it is difficult to obtain the analytic solution of Eq. (3.16), we numerically solve this equation during inflation. Here we assume that the initial amplitude of $Y_+(k, t)$ and that of $Y_-(k, t)$ are the same value and we take the time t_1 as the initial time. In the inflationary stage, the difference between the evolution of $Y_+(k, t)$ and that of $Y_-(k, t)$ is induced by the coupling between the pseudoscalar field and hypercharge electromagnetic fields and thus the hypermagnetic helicity is generated [22, 31, 32, 35]. During inflation ($t_1 \leq t \leq t_R$), the amplitude of $Y_{\pm}(k, t)$ is expressed as

$$Y_{\pm}(k, t) = C_{\pm}(t) Y_{\pm}(k, t_1), \quad (3.17)$$

where $C_{\pm}(t)$ is a numerical value obtained by numerical calculations and we take $C_{\pm}(t_1) = 1$.

In order to obtain the initial amplitude of $Y_{\pm}(k, t)$, we here consider the solution of Eq. (3.16) inside of the horizon, i.e., on subhorizon scales, $k/(aH) \gg 1$. Replacing the independent variable t by η , we find that in the short-wavelength limit, $k \rightarrow \infty$, Eq. (3.16) is approximately given by

$$Y_{\pm}''(k, \eta) + \frac{f'}{f} Y_{\pm}'(k, \eta) + k^2 Y_{\pm}(k, \eta) = 0, \quad (3.18)$$

where the prime denotes differentiation with respect to the conformal time η . Here, in deriving Eq. (3.18), we have taken only the term proportional to k^2 on the right-hand side of Eq. (3.16) and neglected that proportional to k .

The inside solution is given by

$$Y_{\pm}^{\text{in}}(k, \eta) = \frac{1}{\sqrt{2k}} f^{-1/2} e^{-ik\eta}, \quad (3.19)$$

where we have determined the coefficient of this solution by requiring that the vacuum reduces to the one in Minkowski spacetime in the short-wavelength limit. In fact, using Eq. (3.19), we find

$$Y_{\pm}^{\text{in}''}(k, \eta) + \frac{f'}{f} Y_{\pm}^{\text{in}'}(k, \eta) = \left\{ -k^2 - \frac{1}{2} \left[-\frac{1}{2} \left(\frac{f'}{f}\right)^2 + \frac{f''}{f} \right] \right\} Y_{\pm}^{\text{in}}(k, \eta) \quad (3.20)$$

$$\approx -k^2 Y_{\pm}^{\text{in}}(k, \eta), \quad (3.21)$$

where the approximate equality in Eq. (3.21) follows from $-k\eta \gg 1$. Hence we see that the inside solution in Eq. (3.19) approximately satisfies Eq. (3.18).

Here we assume that the initial amplitude of $Y_{\pm}(k, t)$ is approximately given by the inside solution in Eq. (3.19). Hence the initial amplitude of $Y_{\pm}(k, t)$ at the time t_1 is given by

$$|Y_{\pm}(k, t_1)| \approx \frac{1}{\sqrt{2kf(t_1)}}. \quad (3.22)$$

The numerical results of the evolution of $C_{\pm}(t)$ are shown in Figs. 1 and 2. Fig. 1 depicts the case in which $H_{\text{inf}} = 10^{10}\text{GeV}$, $m = 10^9\text{GeV}$, $\lambda = -133$, $\tilde{\lambda} = 1.0$, $w = 1/(75)$, $\phi_1 = 10^9\text{GeV}$, $M = 10^9\text{GeV}$, and $g_{\text{ps}} = 1.0$ (the case (ii) in Table I shown in the next subsection). On the other hand, Fig. 2 depicts the case in which $H_{\text{inf}} = 10^{10}\text{GeV}$, $m = 10^{12}\text{GeV}$, $\lambda = -135$, $\tilde{\lambda} = 1.0$, $w = 1/(75)$, $\phi_1 = 10^{12}\text{GeV}$, $M = 10^{12}\text{GeV}$, and $g_{\text{ps}} = 1.0$ (the case (iii) in Table I). In Figs. 1 and 2, we have used the evolution of ϕ in the positive sign equation in (3.7) and that in Eq. (3.8), respectively. Moreover, we have used $k/a = \exp[-H_{\text{inf}}(t - t_1)]H_{\text{inf}}$, which follows from Eq. (3.1) and $k/(a_1 H_{\text{inf}}) = 1$. The solid curves represents $C_+(t)$ and the dotted curves represents $C_-(t)$. Here we have started to calculate Eq. (3.16) numerically at $t = t_1 = H_{\text{inf}}^{-1}$ and we assume that the initial value of $C_{\pm}(t)$ at $t = t_1$ is $C_{\pm}(t_1) = 1.0$. From Figs. 1 and 2, we understand that after several Hubble expansion times, both the value of $C_+(t)$ and that of $C_-(t)$ becomes almost constant. This qualitative behavior is common to the case $m < H_{\text{inf}}$ and the case $m \gg H_{\text{inf}}$. Finally, we note that if we take a larger value of g_{ps} and/or that of ϕ_1/M (in all the cases in Table I we have taken $\phi_1/M = 1.0$), the difference between the evolution and amplitude of $C_+(t)$ and those of $C_-(t)$ becomes larger.

C. Chern-Simons number density

In this subsection, we consider the generation of the Chern-Simons number density in the inflationary stage and estimate the resultant baryon asymmetry. The proper hyperelectric and hypermagnetic fields are given by [19]

$$E_{Y_i}^{\text{proper}}(t, \mathbf{x}) = a^{-1}E_{Y_i}(t, \mathbf{x}) = -a^{-1}\dot{Y}_i(t, \mathbf{x}), \quad (3.23)$$

$$B_{Y_i}^{\text{proper}}(t, \mathbf{x}) = a^{-1}B_{Y_i}(t, \mathbf{x}) = a^{-2}\epsilon_{ijk}\partial_j Y_k(t, \mathbf{x}), \quad (3.24)$$

where $E_{Y_i}(t, \mathbf{x})$ and $B_{Y_i}(t, \mathbf{x})$ are the comoving hyperelectric and hypermagnetic fields, and ϵ_{ijk} is the totally antisymmetric tensor ($\epsilon_{123} = 1$).

The density of the baryonic number n_{B} is given by [6]

$$n_{\text{B}} = -\frac{n_{\text{f}}}{2}\Delta n_{\text{CS}}, \quad (3.25)$$

$$\Delta n_{\text{CS}} = -\frac{g'^2}{4\pi^2} \int^t \mathbf{E}_Y \cdot \mathbf{B}_Y d\tilde{t}. \quad (3.26)$$

Here, n_{f} is the number of fermionic generations (throughout this paper we use $n_{\text{f}} = 3$) and Δn_{CS} is the Chern-Simons number density.

It follows from Eqs. (3.23) and (3.24) that the Fourier modes $E_{Y_{\pm}}^{\text{proper}}(k, t) = E_{Y_1}^{\text{proper}}(k, t) \pm iE_{Y_2}^{\text{proper}}(k, t)$ and $B_{Y_{\pm}}^{\text{proper}}(k, t) = B_{Y_1}^{\text{proper}}(k, t) \pm iB_{Y_2}^{\text{proper}}(k, t)$ satisfy the

following relations:

$$E_{Y\pm}^{\text{proper}}(k, t) = \pm \frac{1}{k} \frac{\partial B_{Y\pm}^{\text{proper}}(k, t)}{\partial t}, \quad (3.27)$$

$$E_{Y\pm}^{\text{proper}}(k, t) B_{Y\pm}^{\text{proper}}(k, t) = \pm \frac{1}{2} \frac{1}{k} \frac{\partial [B_{Y\pm}^{\text{proper}}(k, t)]^2}{\partial t}. \quad (3.28)$$

On the other hand, the energy density of the proper hypermagnetic field in Fourier space is given by

$$\rho_{B_Y}(k, t) = \frac{1}{2} \left[|B_{Y+}^{\text{proper}}(k, t)|^2 + |B_{Y-}^{\text{proper}}(k, t)|^2 \right] f, \quad (3.29)$$

$$|B_{Y\pm}^{\text{proper}}(k, t)|^2 = \frac{1}{a^2} \left(\frac{k}{a} \right)^2 |Y_{\pm}(k, t)|^2. \quad (3.30)$$

In deriving Eq. (3.30), we have used Eq. (3.24). Multiplying $\rho_{B_Y}(k, t)$ by phase-space density: $4\pi k^3/(2\pi)^3$, we obtain the energy density of the proper magnetic field in the position space

$$\rho_{B_Y}(L, t) = \frac{k^3}{4\pi^2} \left[|B_{Y+}^{\text{proper}}(k, t)|^2 + |B_{Y-}^{\text{proper}}(k, t)|^2 \right] f, \quad (3.31)$$

on a comoving scale $L = 2\pi/k$.

Using Eqs. (3.17), (3.26), (3.28), (3.30), and (3.31), we find that the Chern-Simons number density in the inflationary stage is given by

$$\Delta n_{\text{CS}} = -\frac{g'^2}{4\pi^2} \frac{1}{k} \frac{1}{f} \rho_{B_Y}(L, t) \mathcal{A}(t), \quad (3.32)$$

$$\mathcal{A}(t) = \frac{|C_+(t)|^2 - |C_-(t)|^2}{|C_+(t)|^2 + |C_-(t)|^2}. \quad (3.33)$$

Here we consider the case in which after inflation the Universe is reheated immediately at $t = t_{\text{R}}$. Moreover, we assume that the instantaneous reheating stage is much before the EWPT (the background temperature at the EWPT is $T_{\text{EW}} \sim 100\text{GeV}$). The conductivity of the Universe σ_c is negligibly small during inflation, because there are few charged particles at that time. After reheating, however, a number of charged particles are produced, so that the conductivity immediately jumps to a large value: $\sigma_c \gg H$ ($t \geq t_{\text{R}}$). Consequently, for a large enough conductivity at the instantaneous reheating stage, hyperelectric fields accelerate charged particles and dissipate. On the other hand, the proper hypermagnetic fields evolve in proportion to $a^{-2}(t)$ in the radiation-dominated stage and the subsequent matter-dominated stage ($t \geq t_{\text{R}}$) [19, 24]. Furthermore, the hypermagnetic helicity, i.e., the Chern-Simons number, is conserved [6, 39, 40]. The Chern-Simons number will be released at the EWPT in the form of fermions, which will not be destroyed by the sphaleron processes [14] if the EWPT is strongly first order [5, 6]. Although the EWPT is not strongly first order in the MSM, the EWPT could be strongly first order in the extensions of the MSM [15]. Moreover, at the EWPT the hypermagnetic fields are replaced by the ordinary magnetic fields [5, 6].

Moreover, we discuss the finite conductivity effects. We consider the case in which in the last stage of inflation the conductivity has a finite value. In this case, the term of the conductivity σ_c is added to the inside of the parentheses in the second term on the right-hand side of Eq. (3.16), which is the coefficient of $\dot{Y}_{\pm}(k, t)$. On the other hand, there also exists the term of $\dot{f}/f$ in the inside of the parentheses in the second term on the right-hand side of Eq. (3.16). Hence the term of $\dot{f}/f$ has the same effect as the finite conductivity σ_c . As the physical effect, the term of $\dot{f}/f$ makes both the value of $C_+(t)$ and that of $C_-(t)$ in Eq. (3.17) becomes almost constant after several Hubble expansion times. In fact, in the case in which in the last stage of inflation the conductivity σ_c has a finite value, e.g., $\sigma_c = 100H_{\text{inf}}$, we have numerically calculated the evolution of $C_+(t)$ and $C_-(t)$. As a result, we have found that the results in the case in which we take into account the finite conductivity effects are almost same as the results without taking into account the finite conductivity effects. This is because there exists the term of $\dot{f}/f$ in the coefficient of $\dot{Y}_{\pm}(k, t)$ in Eq. (3.16) and this term has the same physical effect as the finite conductivity σ_c . In the models in Refs. [31, 32], the finite conductivity effects have a influence on results because in these models the coupling $f(\Phi)$ between the dilaton and the hypercharge electromagnetic fields is not considered and hence there does not exist the term of $\dot{f}/f$ in the coefficient of $\dot{Y}_{\pm}(k, t)$. In our model, however, since we consider the coupling $f(\Phi)$, there exists the term of $\dot{f}/f$ in the coefficient of $\dot{Y}_{\pm}(k, t)$ in Eq. (3.16), which has the same physical effect as the finite conductivity σ_c . Consequently, in our model, the finite conductivity effects have little influence on results. Thus we think that the assumption that after reheating the conductivity immediately jumps to a large value, $\sigma_c \gg H$ ($t \geq t_R$), is appropriate, and that the results under this assumption are proper.

It follows from Eqs. (3.25), (3.32), and (3.33) that after the EWPT, the ratio of the density of the baryonic number n_B to the entropy density s is given by

$$\frac{n_B}{s} = n_f \frac{g'^2}{8\pi^2} \frac{1}{k} \frac{\rho_B(L, t)}{s} \mathcal{A}(t_R), \quad (3.34)$$

with

$$\rho_B(L, t) = \frac{1}{8\pi^2} \frac{1}{f(t_1)} \left(\frac{k}{a} \right)^4 [|C_+(t_R)|^2 + |C_-(t_R)|^2], \quad (3.35)$$

$$f(t_1) = \left(1 - \tilde{\lambda}^2 w N \right)^{-X}, \quad (3.36)$$

$$N = H_{\text{inf}} (t_R - t_1), \quad (3.37)$$

$$X \equiv \frac{\lambda}{\tilde{\lambda}}, \quad (3.38)$$

where we have used the fact that when $t \geq t_R$, $f = 1$, and that after the EWPT, $\rho_{B_Y}(L, t) \rightarrow \rho_B(L, t)$, where $\rho_B(L, t)$ is the energy density of the ordinary magnetic fields. Moreover, we have taken into account the fact that for a large enough conductivity after the instantaneous reheating stage, the proper hypermagnetic fields evolve in proportion to $a^{-2}(t)$ as stated above. Here, N is the number of e -folds between the time t_1 and the end of inflation t_R . Furthermore, in deriving Eq. (3.36), we have used Eqs. (2.6) and (3.4) with $\Phi_R = 0$.

TABLE I: Estimates of the value of n_B/s for $w = 1/(75)$, $\phi_1 = M = m$, $g_{\text{ps}} = 1.0$, and $\tilde{\lambda} = 1.0$. Here we have used the evolution of ϕ in the positive sign equation in (3.7) for the cases (i), (ii), and (iv), and that in Eq. (3.8) for those (iii) and (v).

	$ n_B/s $	$B(H_0^{-1}, t_0)$ [G]	H_{inf} [GeV]	m [GeV]	$C_+(t_R)$	$C_-(t_R)$	X
(i)	3.6×10^{-10}	2.7×10^{-24}	1.0×10^{14}	1.0×10^{12}	0.367	2.23	-1.11×10^2
(ii)	1.0×10^{-10}	1.4×10^{-24}	1.0×10^{10}	1.0×10^9	0.369	2.18	-1.33×10^2
(iii)	1.5×10^{-10}	3.5×10^{-24}	1.0×10^{10}	1.0×10^{12}	1.04	0.816	-1.35×10^2
(iv)	0.96×10^{-10}	1.4×10^{-24}	1.0×10^5	1.0×10^3	0.368	2.17	-1.65×10^2
(v)	2.8×10^{-10}	4.7×10^{-24}	1.0×10^5	1.0×10^7	1.04	0.808	-1.68×10^2

In order to estimate the value of n_B/s , we use the following relations [18]:

$$H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1} = 2.1h \times 10^{-42} \text{ GeV}, \quad (3.39)$$

$$\rho_\varphi(t_R) = \frac{\pi^2}{30} g_R T_R^4 \quad (g_R \approx 200), \quad (3.40)$$

$$N = 45 + \ln \left(\frac{L}{[\text{Mpc}]} \right) + \ln \left\{ \frac{[30/(\pi^2 g_R)]^{1/12} \rho_\varphi^{1/4}}{10^{38/3} [\text{GeV}]} \right\}, \quad (3.41)$$

$$\frac{a_R}{a_0} = \left(\frac{g_R}{3.91} \right)^{-1/3} \frac{T_{\gamma 0}}{T_R} \approx \frac{2.35 \times 10^{-13} [\text{GeV}]}{3.71 T_R} \quad (T_{\gamma 0} \approx 2.73 \text{ K}), \quad (3.42)$$

$$s_0 = 2.97 \times 10^3 \left(\frac{T_{\gamma 0}}{2.75 [\text{K}]} \right)^3 \text{ cm}^{-3}, \quad (3.43)$$

where H_0 is the Hubble constant at the present time (throughout this paper we use $h = 0.70$ [41]), g_R is the total number of degrees of freedom for relativistic particles at the reheating epoch, T_R is reheating temperature, $T_{\gamma 0}$ is the present temperature of the cosmic microwave background (CMB) radiation, and s_0 is the entropy density at the present time. Moreover, we use $g'^2/(4\pi) = \alpha_{\text{EM}}/\cos^2 \theta_w$, where $\alpha_{\text{EM}} = 1/(137)$ is the fine-structure constant and θ_w is the weak mixing angle. The experimentally measured value of θ_w is given by $\sin^2 \theta_w \simeq 0.23$ [12].

Consequently, from Eq. (3.34), we can estimate the value of the ratio of the density of the baryonic number n_B to the entropy density s , which is observationally estimated as $n_B/s = 0.92 \times 10^{-10}$ by using the the first year Wilkinson Microwave Anisotropy Probe (WMAP) data on the anisotropy of the CMB radiation [42]. Table I displays the estimate of the value of n_B/s for $w = 1/(75)$, $\phi_1 = M = m$, $g_{\text{ps}} = 1.0$, and $\tilde{\lambda} = 1.0$. Here we have used the evolution of ϕ in the positive sign equation in (3.7) for the cases (i), (ii), and (iv), and that in Eq. (3.8) for those (iii) and (v). In Table I, we have considered the BAU induced by the helicity of the hypermagnetic fields whose present scale is the present horizon scale, $L = H_0^{-1}$. Hence the resultant BAU is homogeneous over the present horizon. Moreover, in estimating the value of the energy density of the magnetic fields in Eq. (3.35), we have used $a_R/a_1 = \exp(N)$ and $k/(a_1 H_{\text{inf}}) = 1$. Furthermore, a constraint on H_{inf} from tensor

perturbations [43, 44] is obtained by using the WMAP three year data on temperature fluctuations [45], $H_{\text{inf}} < 5.9 \times 10^{14} \text{GeV}$.

From Table I, we see that if the magnetic fields with the field strength $\sim 10^{-24} \text{G}$ on the horizon scale at the present time are generated, the resultant value of n_{B}/s can be as large as 10^{-10} , which is consistent with the above observational estimation of WMAP. When the generated magnetic fields with the field strength $\sim 10^{-24} \text{G}$ on the horizon scale at the present time play the role of seed magnetic fields of galactic magnetic fields, by assuming the magnetic flux conservation, $Br^2 = \text{constant}$, where r is a scale, and using the scale ratio $r_{\text{gal}}/H_0^{-1} \sim 10^{-5}$, where r_{gal} is the scale of galaxies, we can estimate the seed magnetic field strength of the galactic magnetic fields as $B_{\text{gal}}^{(\text{seed})} \sim 10^{-14} \text{G}$ which is sufficient for the following dynamo amplification. Thus, in this way, the main problem of cosmological magnetic fields can be solved by obtaining both large spatial scale and large magnetic field amplitude as it should be for the relic seed magnetic fields obeying dynamo theories.

Here we state the reason why in this model the amplitude of the generated magnetic fields can be as large as $\sim 10^{-24} \text{G}$ on the horizon scale at the present time. The reason is that the conformal invariance of the hypercharge electromagnetic fields is extremely broken through the coupling between the dilaton and the hypercharge electromagnetic fields by introducing a huge hierarchy between the coupling constant of the dilaton to the hypercharge electromagnetic fields λ and the coupling constant $\tilde{\lambda}$ of the dilaton potential, $|X| = |\lambda/\tilde{\lambda}| \gg 1$. From Table I, we see that in order that the magnetic fields with the field strength $\sim 10^{-24} \text{G}$ on the horizon scale at the present time could be generated, we have to introduce the huge hierarchy $|X| \gg 1$. It follows from Eqs. (3.22) and (3.36) that if $|X| \gg 1$, the value of $f(t_1)$ is very small and hence the amplitude of $|Y_{\pm}(k, t_1)|$ is sufficiently large. Moreover, it follows from Figs. 1 and 2 that after several Hubble expansion times, both the value of $C_+(t)$ and that of $C_-(t)$ becomes almost constant. Hence it follows from Eq. (3.17) that the amplitude of $|Y_{\pm}(k, t)|$ is sufficiently large. Consequently, it follows from Eqs. (3.30), (3.31), and (3.35) that the energy density of the generated magnetic fields, i.e., the resultant magnetic field amplitude can be as large as $\sim 10^{-24} \text{G}$ on the horizon scale at the present time. This is the reason why the present amplitude of the generated magnetic fields in this model is larger than that in other inflation scenarios [20].

The physical reason why the coupling of the dilaton to the hypercharge electromagnetic fields has to be much stronger than the coupling in the dilaton potential in order that the amplitude of the generated magnetic fields could be as large as $\sim 10^{-24} \text{G}$ on the horizon scale at the present time is as follows: In order to generate the magnetic fields with the sufficient strength, the conformal invariance of the hypercharge electromagnetic fields has to be extremely broken through the coupling $f(\Phi)$ between the dilaton and the hypercharge electromagnetic fields. In order that this condition could be met, it is necessary that the change of the value of $f(\Phi)$ in the inflationary stage is much larger than the change of the value of the dilaton potential $V[\Phi]$ at that stage, in other words, $f(\Phi)$ evolves more rapidly than the dilaton potential $V[\Phi]$. Thus the coupling of the dilaton to the hypercharge electromagnetic fields has to be much stronger than the coupling in the dilaton potential, i.e., the value of $|\lambda|$ has to be much larger than that of $\tilde{\lambda}$. In this case, the value of $f(\Phi)$ can change from $f(t_1) \ll 1$ to $f(t_{\text{R}}) = 1$ in the inflationary stage ($t_1 \leq t \leq t_{\text{R}}$). Consequently, from Eqs. (3.35) and (3.36), we see that the amplitude of the generated magnetic fields can be sufficiently large because $f(t_1)$ is much smaller than unity.

IV. CONCLUSION

In the present paper we have studied the generation of the BAU from the helicity of hypermagnetic fields in inflationary cosmology, taking into account the breaking of the conformal invariance of the hypercharge electromagnetic fields by introducing both a coupling with the dilaton and that with a pseudoscalar field. Owing to the coupling between the pseudoscalar field and the hypercharge electromagnetic fields, the hypermagnetic helicity, which corresponds to the Chern-Simons number, is generated in the inflationary stage. The Chern-Simons number stored in the hypercharge electromagnetic fields is converted into fermions at the EWPT due to the Abelian anomaly, and at the same time the hypermagnetic fields are replaced by the ordinary magnetic fields, which survive after the EWPT. The generated fermions can survive after the EWPT if the EWPT is strongly first order. In the extensions of the MSM, the above condition could be satisfied.

As a result, we have found that if the magnetic fields with sufficient strength on the horizon scale at the present time are generated, the resultant value of the ratio of the density of the baryonic number to the entropy density, n_B/s , can be as large as 10^{-10} , which is consistent with the magnitude of the BAU suggested by observations obtained from WMAP.

Acknowledgements

The author is deeply grateful to Mikhail Shaposhnikov, Jun'ichi Yokoyama, and Yasunari Kurita for helpful discussions. The author's work was partially supported by the Monbu-Kagaku Sho 21st century COE Program "Center for Diversity and Universality in Physics" and was also supported by a Grant-in-Aid provided by the Japan Society for the Promotion of Science.

-
- [1] A. D. Sakharov, JETP Lett. **5**, 24 (1967).
 - [2] A. D. Dolgov, Phys. Rep. **222**, 309 (1992).
 - [3] M. Dine, A. Kusenko, Rev. Mod. Phys. **76**, 1 (2004).
 - [4] M. Joyce and M. Shaposhnikov, Phys. Rev. Lett. **79**, 1193 (1997).
 - [5] M. Giovannini and M. E. Shaposhnikov, Phys. Rev. Lett. **80**, 22 (1998).
 - [6] M. Giovannini and M. E. Shaposhnikov, Phys. Rev. D **57**, 2186 (1998).
 - [7] C. Thompson, Phys. Lett. B **422**, 61 (1998).
 - [8] T. Vachaspati and G. B. Field, Phys. Rev. Lett. **73**, 373 (1994).
 - [9] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Ann. Rev. Nucl. Part. Sci. **43**, 27 (1993).
 - [10] V. A. Rubakov and M. E. Shaposhnikov, Usp. Fiz. Nauk **166**, 493 (1996) [Phys. Usp. **39**, 461 (1996)].
 - [11] K. Funakubo, Prog. Theor. Phys. **96**, 475 (1996).
 - [12] M. Trodden, Rev. Mod. Phys. **71**, 1463 (1999).
 - [13] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).
 - [14] N. S. Manton, Phys. Rev. D **28**, 2019 (1983); F. R. Klinkhamer and N. S. Manton, *ibid.* **30**, 2212 (1984).
 - [15] K. Funakubo, S. Tao, and F. Toyoda, Prog. Theor. Phys. **114**, 369 (2005).

- [16] M. S. Turner and L. M. Widrow, Phys. Rev. D **37**, 2743 (1988).
- [17] A. D. Linde, *Particle Physics and Inflationary Cosmology* (Harwood Academic, Chur, Switzerland, 1990); K. A. Olive, Phys. Rep. **190**, 307 (1990); D. H. Lyth and A. Riotto, *ibid.* **314**, 1, (1999).
- [18] E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley, Redwood City, California, 1990).
- [19] B. Ratra, Astrophys. J. **391**, L1 (1992); Report No. GRP-287/CALT-68-1751.
- [20] D. Lemoine and M. Lemoine, Phys. Rev. D **52**, 1955 (1995); M. Gasperini, M. Giovannini, and G. Veneziano, Phys. Rev. Lett. **75**, 3796 (1995).
- [21] W. D. Garretson, G. B. Field, and S. M. Carroll, Phys. Rev. D **46**, 5346 (1992).
- [22] G. B. Field and S. M. Carroll, Phys. Rev. D **62**, 103008 (2000).
- [23] A. D. Dolgov, Phys. Rev. D **48**, 2499 (1993).
- [24] K. Bamba and J. Yokoyama, Phys. Rev. D **69**, 043507 (2004).
- [25] K. Bamba and J. Yokoyama, Phys. Rev. D **70**, 083508 (2004).
- [26] D. Grasso and H. R. Rubinstein, Phys. Rep. **348**, 163 (2001).
- [27] A. D. Dolgov, hep-ph/0110293.
- [28] L. M. Widrow, Rev. Mod. Phys. **74**, 775 (2002).
- [29] M. Giovannini, Int. J. Mod. Phys. D **13**, 391 (2004).
- [30] V. B. Semikoz and D. D. Sokoloff, Int. J. Mod. Phys. D **14**, 1839 (2005).
- [31] R. Brustein and D. H. Oaknin, Phys. Rev. Lett. **82**, 2628 (1999); Phys. Rev. D **60**, 023508 (1999).
- [32] M. Giovannini, Phys. Rev. D **61**, 063004 (2000); **61**, 063502 (2000).
- [33] R. Brustein and D. H. Oaknin, Phys. Rev. D **63**, 055002 (2001) ;Phys. Rev. D **62**, 015001 (2000).
- [34] E. I. Guendelman and D. A. Owen, Phys. Lett. B **276**, 108 (1992).
- [35] L. Campanelli and M. Giannotti, Phys. Rev. D **72**, 123001 (2005).
- [36] J. P. Derendinger, L. E. Ibáñez, and H. P. Nilles, Phys. Lett. **155B**, 65 (1985); M. Dine, R. Rohm, N. Seiberg, and E. Witten, *ibid.* **156B**, 55 (1985).
- [37] T. Barreiro, B. de Carlos, and E. J. Copeland, Phys. Rev. D **58**, 083513 (1998).
- [38] A. D. Dolgov, K. Kohri, O. Seto, and J. Yokoyama, Phys. Rev. D **67**, 103515 (2003).
- [39] E. N. Parker, Astrophys. J. **163**, 255 (1971); *Cosmical Magnetic Fields* (Clarendon, Oxford, England, 1979); Ya. B. Zel'dovich, A. A. Ruzmaikin, and D. D. Sokoloff, *Magnetic Fields in Astrophysics* (Gordon and Breach, New York, 1983).
- [40] D. Biskamp, *Nonlinear Magnetohydrodynamics* (Cambridge University Press, Cambridge, United Kingdom, 1993).
- [41] W. L. Freedman *et al.*, Astrophys. J. **553**, 47 (2001).
- [42] D. N. Spergel *et al.*, Astrophys. J., Suppl. Ser. **148**, 175 (2003).
- [43] V. A. Rubakov, M. V. Sazhin, and A. V. Veryaskin, Phys. Lett. **115B**, 189 (1982).
- [44] L. F. Abbott and M. B. Wise, Nucl. Phys. **B244**, 541 (1984).
- [45] D. N. Spergel *et al.*, astro-ph/0603449.




FIG. 1: Numerical results of the evolution of $C_{\pm}(t)$ for $H_{\text{inf}} = 10^{10}\text{GeV}$, $m = 10^9\text{GeV}$, $\lambda = -133$, $\tilde{\lambda} = 1.0$, $w = 1/(75)$, $\phi_1 = 10^9\text{GeV}$, $M = 10^9\text{GeV}$, and $g_{\text{ps}} = 1.0$ (the case (ii) in Table I). Here we have used the evolution of ϕ in the positive sign equation in (3.7). The solid curves represents $C_+(t)$ and the dotted curves represents $C_-(t)$.




FIG. 2: Numerical results of the evolution of $C_{\pm}(t)$ for $H_{\text{inf}} = 10^{10}\text{GeV}$, $m = 10^{12}\text{GeV}$, $\lambda = -135$, $\tilde{\lambda} = 1.0$, $w = 1/(75)$, $\phi_1 = 10^{12}\text{GeV}$, $M = 10^{12}\text{GeV}$, and $g_{\text{ps}} = 1.0$ (the case (iii) in Table I). Here we have used the evolution of ϕ in Eq. (3.8). The solid curves represents $C_+(t)$ and the dotted curves represents $C_-(t)$.